North Carolina Department of Health and Human Services Women's and Children's Health Section • Immunization Branch

MEDICAL EXEMPTION STATEMENT

Purpose: To provide physicians, licensed to practice medicine in North Carolina, a mechanism to document a true medical contraindication/precaution to an immunization(s). This form does not need approval from the State Health Director. This form can be accepted by agencies that require proof of immunizations. For medical exemptions NOT listed in the table below, submit the Physician's Request for Medical Exemption form (Form: DHHS 3995) to the State Health Director for approval, available at

http://www.immunize.nc.gov/schools/ncexemptions.htm

Name of Patient	DOB
Name of Parent/Guardian	Primary Phone ()
Home Address (Patient/Parent)	County
Name of Child Care/School/College/University	·

Medical contraindications and precautions for immunizations are described in the most recent recommendations of the Advisory Committee on Immunization Practices (ACIP), available at http://www.cdc.gov/vaccines/recs/vac-admin/contraindications.htm

A contraindication is a condition in a recipient that increases the risk for a serious adverse reaction. A vaccine will not be administered when a contraindication is present. A precaution is a condition in a recipient that might increase the risk for a serious adverse reaction or that might compromise the ability of the vaccine to produce immunity. Under normal conditions, vaccinations should be deferred when a precaution is present.

,			
	Vaccine	.+	Check all true contraindications and precautions that apply to this patient below:
	Diphtheria,		Contraindications
	tetanus,		Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component.
	pertussis		For pertussis-containing vaccines: encephalopathy (e.g., coma, decreased level of consciousness, prolonged
	(DТаР)		seizure) not attributable to another identifiable cause within 7 days of administration of DTaP or DTP (for DTaP); or of previous dose of DTaP, DTP, or Tdap (for Tdap).
·Ü	Tetanus,		Precautions
	diphtheria,		Moderate or severe acute illness with or without fever.
	pertussis		Guillain-Barré syndrome (GBS) within 6 weeks after a previous dose of tetanus toxoid-containing vaccine.
	(Tdap)		History of arthus-type hypersensitivity reaction after a previous dose of a tetanus or diphtheria toxoid-containing vaccine; defer until at least 10 years have elapsed since the last tetanus-toxoid containing vaccine.
0	Tetanus, diphtheria (DT, Td)	0	For pertussis-containing vaccines: progressive or unstable neurologic disorder (including infantile spasms for DTaP), uncontrolled seizures, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized.
			Additional Precautions that only apply to DTaP
			Temperature of 105° For higher (40.5° Cor higher) within 48 hours after vaccination with a previous dose of DTP/DTaP.
		D	Collapse or shock-like state (i.e., hypotonic hyporesponsive episode) within 48 hours after receiving a previous dose of DTP/DTaP.
			Seizure within 3 days after receiving a previous dose of DTP/DTaP.
		· 🛮	Persistent, inconsolable crying lasting 3 or more hours within 48 hours after receiving a previous dose of DTP/DTaP.
	Measles,		Contraindications
	mumps,	Ū	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component.
	rubella	П	Known severe immunodeficiency (e.g., congenital immunodeficiency, malignancy, chemotherapy, long-term
	(MMR)		immunosuppressive therapy, or human immunodeficiency virus [HIV] infection with CD4+ T-lymphocyte count ≤
			15%).
			Pregnancy.
			Precautions
			Moderate or severe acute Illness with or without fever.
			Recent (within 11 months) receipt of antibody-containing blood product (specific interval depends on product).
	1	О	History of thrombocytopenia or thrombocytopenic purpuca.
			Need for tuberculin skin testing (Measles vaccine might suppress tuberculin reactivity temporarily)

 □ Varicella	Contraindications
(Var)	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component. Known severe immunodeficiency (e.g., congenital immunodeficiency, malling new characteristics)
(001)	- Management (e.g., confermed intributorenciette), manginancy, chelifotherapy, long-term
1	immunosuppressive therapy, or human immunodeficiency virus (HIV) infection with CD4+ T-lymphocyte count ≤ 15%.
	□ Pregnancy.
	Precautions
	Moderate or severe acute illness with or without fever
	Recent (within 11 months) receipt of antibody-containing blood product (specific interval depends on product) Receipt of specific antivirals (e.g., acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination, Avoid use of
	these antivirals for 14 days after vaccination.
☐ Inactivated	Contraindications
Polio Virus	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component.
(IPV)	Precautions
·	Moderate or severe acute Illness with or without fever.
	□ Pregnancy.
☐ Hepatitis B	Contraindications
(Hep B)	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component.
	Precautions
	☐ Moderate or severe acute illness with or without fever.
	Infant weighing less than 2000 grams (4 lbs, 6.4 oz) if mother is documented hepatitis B surface antigen (HbsAg)
	negative at the time of the infant's birth.
☐ Haemophīlus	Contraindications
<i>Influenza</i> tγpe B	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component.
(HIB)	☐ Age younger than 6 weeks. Precautions
(unit	Moderate or severe acute illness with or without fevery
Pneumococcal	Contraindications
(PCV13)	Severe allergic reaction (e.g., anaphylaxis) after a previous dose, or to a vaccine component, including any
	diphtheria toxoid-containing vaccine.
	Precautions
	Moderate or severe acute illness with or without fever.
☐ Meningococcal	Contraindications
(MCV4)	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	Precautions
	☐ Moderate or severe acute illness with or without fever
A physicia	in (M.D. or D.O) licensed to practice medicine in North Carolina must complete and sign this form.
	ds:
	·
N.C. Physician's Nan	ne (please print) Phone
	atureDate
	Instructions:
1. Complete and sign	
2. Attach a copy of t	he most current immunization record,
	the patient's medical record.
	to the person requesting this form.
	· p

For questions call (919) 707-5550

Additional copies of this form can be accessed at: http://www.immunize.nc.gov/schools/ncexemptions.htm

DHHS 3987 (Revised 6/15) Immunization (Review:6/17)